



An Equivalent to the Riemann Hypothesis

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

December 19, 2023

An Equivalent to the Riemann Hypothesis

Frank Vega

To my mother

Abstract. The Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. It is considered by many to be the most important unsolved problem in pure mathematics. There are several statements equivalent to the famous Riemann hypothesis. Robin's criterion states that the Riemann hypothesis is true if and only if the inequality $\sigma(n) < e^\gamma \cdot n \cdot \log \log n$ holds for all natural numbers $n > 5040$, where $\sigma(n)$ is the sum-of-divisors function of n , $\gamma \approx 0.57721$ is the Euler-Mascheroni constant and \log is the natural logarithm. We prove that the Riemann hypothesis is true whenever there exists a large enough positive number x_0 such that for all $x > x_0$ we obtain that the value of

$$\sum_{n \leq \alpha_x} \frac{1}{n} - \sum_{6 \leq n \leq \frac{x}{\log x}} \frac{e^{-\gamma}}{n \cdot (\log(n \cdot \log n))} - \sum_{n < 6} \frac{e^{-\gamma}}{q_n}$$

is lesser than or equal to $e^{-\gamma} \cdot \left(\gamma - B - \frac{1}{2 \cdot (x-1)} \right)$ where $B \approx 0.26149$ is the Meissel-Mertens constant and $\alpha_x = \left(\log x + \frac{0.0222 \cdot \log x}{\log \log x} \right)$. Since the previous expression goes to 0 as x tends to infinity, then we deduce that the Riemann hypothesis must be true.

Mathematics Subject Classification (2010). Primary 11M26; Secondary 11A41, 11A25.

Keywords. Riemann hypothesis, prime numbers, Robin's criterion, superabundant numbers.

1. Introduction

In mathematics, the Prime-counting function $\pi(x)$ is given by

$$\pi(x) = \sum_{q \leq x} 1$$

with the sum extending over all prime numbers q that are less than or equal to x .

Proposition 1.1. For $x \geq 17$ [11, Corollary 1 (3.5) pp. 69]:

$$\pi(x) > \frac{x}{\log x},$$

where \log is the natural logarithm.

Proposition 1.2. For the n th prime number q_n and $n \geq 6$ [11, Corollary (3.13) pp. 69]:

$$q_n < n \cdot (\log(n \cdot \log n)).$$

The number $\gamma \approx 0.57721$ is the Euler-Mascheroni constant which is defined as

$$\begin{aligned} \gamma &= \lim_{n \rightarrow \infty} \left(-\log n + \sum_{k=1}^n \frac{1}{k} \right) \\ &= \int_1^{\infty} \left(-\frac{1}{x} + \frac{1}{[x]} \right) dx. \end{aligned}$$

Here, $[\dots]$ represents the floor function. Franz Mertens discovered some important results about the constants B and H (1874) [5]. We define $H = \gamma - B$ such that $B \approx 0.26149$ is the Meissel-Mertens constant [5].

Proposition 1.3. We have [2, Lemma 2.1 (1) pp. 359]:

$$\sum_{k=1}^{\infty} \left(\log \left(\frac{q_k}{q_k - 1} \right) - \frac{1}{q_k} \right) = \gamma - B = H,$$

where q_k is the k th prime number.

For $x \geq 2$, the function $u(x)$ is defined as follows [7, pp. 379]:

$$u(x) = \sum_{q > x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Proposition 1.4. We have [7, (11) pp. 379]:

$$0 < u(x) \leq \frac{1}{2 \cdot (x-1)}.$$

On the sum of the reciprocals of all prime numbers not exceeding x , we have:

Proposition 1.5. For $x \geq 2278383$ [3, Theorem 5.6 (1) pp. 243]:

$$-\frac{0.2}{\log^3 x} \leq \sum_{q \leq x} \frac{1}{q} - B - \log \log x \leq \frac{0.2}{\log^3 x}$$

As usual $\sigma(n)$ is the sum-of-divisors function of n

$$\sum_{d|n} d,$$

where $d | n$ means the integer d divides n . Define $f(n)$ as $\frac{\sigma(n)}{n}$.

Proposition 1.6. Let $\prod_{i=1}^r q_i^{a_i}$ be the representation of n as a product of prime numbers $q_1 < \dots < q_r$ with natural numbers a_1, \dots, a_r as exponents. Then [4, Lemma 1 pp. 2],

$$f(n) = \left(\prod_{i=1}^r \frac{q_i}{q_i - 1} \right) \cdot \prod_{i=1}^r \left(1 - \frac{1}{q_i^{a_i+1}} \right).$$

Definition 1.7. We say that $\text{Robin}(n)$ holds provided that

$$f(n) < e^\gamma \cdot \log \log n.$$

The Ramanujan's Theorem stated that if the Riemann hypothesis is true, then the previous inequality holds for large enough n . Next, we have the Robin's Theorem:

Proposition 1.8. $\text{Robin}(n)$ holds for all natural numbers $n > 5040$ if and only if the Riemann hypothesis is true [10, Theorem 1 pp. 188].

Unconditionally on Riemann hypothesis, we have:

Proposition 1.9. $\text{Robin}(n)$ holds for all natural numbers $10^{10^{13.11485}} \geq n > 5040$ [9, Theorem 5 pp. 6].

Proposition 1.10. For $x \geq 1$ [11, Corollary 2 (3.31) pp. 71]:

$$\prod_{q \leq x} \frac{q}{q-1} < e^\gamma \cdot \sum_{m \leq x} \frac{1}{m},$$

where m denotes a natural number.

In 1997, Ramanujan's old notes were published where he defined the generalized highly composite numbers, which include the superabundant and colossally abundant numbers [8]. Superabundant numbers were also studied by Leonidas Alaoglu and Paul Erdős (1944) [1]. Let $q_1 = 2, q_2 = 3, \dots, q_k$ denote the first k consecutive primes, then an integer of the form $\prod_{i=1}^k q_i^{a_i}$ with $a_1 \geq a_2 \geq \dots \geq a_k \geq 1$ is called a Hardy-Ramanujan integer [2, pp. 367]. A natural number n is called superabundant precisely when, for all natural numbers $m < n$

$$f(m) < f(n).$$

We know the following property for the superabundant numbers:

Proposition 1.11. If n is superabundant, then n is a Hardy-Ramanujan integer [1, Theorem 1 pp. 450].

Proposition 1.12. Let n be a large enough superabundant number such that q is the largest prime factor of n . Then [6, Corollary 4.16 pp. 16]:

$$q < (\log n) \cdot \left(1 + \frac{0.0222}{\log \log n} \right).$$

A number n is said to be colossally abundant if, for some $\epsilon > 0$,

$$\frac{\sigma(n)}{n^{1+\epsilon}} \geq \frac{\sigma(m)}{m^{1+\epsilon}} \text{ for } (m > 1).$$

There is a close relation between the superabundant and colossally abundant numbers.

Proposition 1.13. *Every colossally abundant number is superabundant [1, pp. 455].*

Several analogues of the Riemann hypothesis have already been proved. Many authors expect (or at least hope) that it is true. However, there are some implications in case of the Riemann hypothesis could be false.

Proposition 1.14. *If the Riemann hypothesis is false, then there exist infinitely many colossally abundant numbers $n > 5040$ such that Robin(n) fails (i.e. Robin(n) does not hold) [10, Proposition pp. 204].*

Putting all together yields a proof for the Riemann hypothesis.

2. Central Lemma

The following is a key Lemma.

Lemma 2.1. *If the Riemann hypothesis is false, then there exist infinitely many superabundant numbers n such that Robin(n) fails.*

Proof. This is a direct consequence of Propositions 1.8, 1.13 and 1.14. \square

3. Main Insight

This is the main insight.

Theorem 3.1. *The Riemann hypothesis is true whenever there exists a large enough positive number n_0 such that for all $n > n_0$ we obtain that the value of*

$$\sum_{m \leq \alpha_n} \frac{1}{m} - \sum_{6 \leq m \leq \frac{n}{\log n}} \frac{e^{-\gamma}}{m \cdot (\log(m \cdot \log m))} - \sum_{m < 6} \frac{e^{-\gamma}}{q_m}$$

is lesser than or equal to $e^{-\gamma} \cdot \left(H - \frac{1}{2 \cdot (n-1)} \right)$ where $\alpha_n = \left(\log n + \frac{0.0222 \cdot \log n}{\log \log n} \right)$.

Proof. Let $n > 5040$ be a counterexample such that Robin(n) does not hold. We know this number could be a large enough superabundant number by Lemma 2.1. Let $\prod_{i=1}^k q_i^{a_i}$ be the representation of this superabundant number n as the product of the first k consecutive primes $q_1 < \dots < q_k$ with the natural numbers $a_1 \geq a_2 \geq \dots \geq a_k \geq 1$ as exponents according to Proposition 1.11. We know that $n > 10^{10^{13.11485}}$ by Proposition 1.9. Under our supposition, we have

$$\sigma(n) \geq e^{\gamma} \cdot n \cdot \log \log n.$$

By Proposition 1.5, we notice that

$$\sum_{q \leq n} \frac{1}{q} \leq \log \log n + B + \frac{0.2}{\log^3 n}.$$

We can see that

$$\sum_{q \leq n} \log \left(\frac{q}{q-1} \right) - \sum_{q \leq n} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right) \leq \log \log n + B + \frac{0.2}{\log^3 n}$$

which is

$$\sum_{q \leq n} \log \left(\frac{q}{q-1} \right) - H < \log \log n + B + \frac{0.2}{\log^3 n}$$

and

$$\sum_{q \leq n} \log \left(\frac{q}{q-1} \right) < \log \log n + \gamma + \frac{0.2}{\log^3 n}$$

by Proposition 1.3. That is the same as

$$n \cdot \sum_{q \leq n} \log \left(\frac{q}{q-1} \right) < n \cdot \log \log n + \left(\gamma + \frac{0.2}{\log^3 n} \right) \cdot n$$

after multiplying both sides by the superabundant number n . We know that

$$\left(\gamma + \frac{0.2}{\log^3 n} \right) \cdot n \leq (e^\gamma - 1) \cdot n \cdot \log \log n$$

for $n > 10^{10^{13.11485}}$. Consequently, we obtain that

$$\sigma(n) > n \cdot \sum_{q \leq n} \log \left(\frac{q}{q-1} \right)$$

by transitivity since

$$e^\gamma \cdot n \cdot \log \log n \geq n \cdot \log \log n + \left(\gamma + \frac{0.2}{\log^3 n} \right) \cdot n.$$

In this way, we have

$$f(n) > \sum_{q \leq n} \log \left(\frac{q}{q-1} \right)$$

which is

$$\prod_{q \leq \alpha_n} \left(\frac{q}{q-1} \right) > \sum_{q \leq n} \log \left(\frac{q}{q-1} \right)$$

by Proposition 1.6 and 1.12 since

$$\prod_{q \leq q_k} \left(\frac{q}{q-1} \right) > f(n)$$

and

$$q_k < (\log n) \cdot \left(1 + \frac{0.0222}{\log \log n} \right) = \left(\log n + \frac{0.0222 \cdot \log n}{\log \log n} \right) = \alpha_n$$

for large enough superabundant number n . That would be

$$e^\gamma \cdot \sum_{m \leq \alpha_n} \frac{1}{m} > \sum_{q \leq n} \log \left(\frac{q}{q-1} \right)$$

since

$$\prod_{q \leq \alpha_n} \left(\frac{q}{q-1} \right) < e^\gamma \cdot \sum_{m \leq \alpha_n} \frac{1}{m}$$

by Proposition 1.10. So, we would have

$$e^\gamma \cdot \sum_{m \leq \alpha_n} \frac{1}{m} - \sum_{q \leq n} \frac{1}{q} > H - u(n)$$

by Proposition 1.3. By Proposition 1.4, we have

$$e^\gamma \cdot \sum_{m \leq \alpha_n} \frac{1}{m} - \sum_{q \leq n} \frac{1}{q} > H - \frac{1}{2 \cdot (n-1)}.$$

That is equivalent to

$$e^\gamma \cdot \sum_{m \leq \alpha_n} \frac{1}{m} - \sum_{m \leq \frac{n}{\log n}} \frac{1}{q_m} > H - \frac{1}{2 \cdot (n-1)}$$

since

$$\pi(n) > \frac{n}{\log n}$$

by Proposition 1.1. Hence, it is enough to show that

$$\sum_{m \leq \alpha_n} \frac{1}{m} - \sum_{m \leq \frac{n}{\log n}} \frac{e^{-\gamma}}{q_m} > e^{-\gamma} \cdot \left(H - \frac{1}{2 \cdot (n-1)} \right)$$

which means that the expression

$$\sum_{m \leq \alpha_n} \frac{1}{m} - \sum_{6 \leq m \leq \frac{n}{\log n}} \frac{e^{-\gamma}}{m \cdot (\log(m \cdot \log m))} - \sum_{m < 6} \frac{e^{-\gamma}}{q_m}$$

would be greater than $e^{-\gamma} \cdot \left(H - \frac{1}{2 \cdot (n-1)} \right)$ by Proposition 1.2. However, that contradicts the fact that n could be a superabundant number as large as we want and thus, it could happen that $n > n_0$ from our pre-conditions. This contradiction implies that it cannot exist infinitely many superabundant numbers n such that $\text{Robin}(n)$ fails and therefore, the Riemann hypothesis should be true using a proof by contraposition from Lemma 2.1. \square

4. Main Theorem

This is the main theorem.

Theorem 4.1. *The Riemann hypothesis is true.*

Proof. Since the expression of Theorem 3.1 goes to 0 as n tends to infinity, then we deduce that the Riemann hypothesis must be true. This is because of the upper bound $\frac{n}{\log n}$ is exponentially larger than the number $\alpha_n = \left(\log n + \frac{0.0222 \cdot \log n}{\log \log n} \right)$ and

$$\frac{e^{-\gamma}}{m \cdot (\log(m \cdot \log m))} \approx \frac{1}{m}$$

is true for each value of $m \geq 6$ since $e^\gamma \approx 0$ according to the Proposition 1.2. \square

References

- [1] Alaoglu, L., Erdős, P.: On Highly Composite and Similar Numbers. *Transactions of the American Mathematical Society* **56**(3), 448–469 (1944). <https://doi.org/10.2307/1990319>
- [2] Choie, Y., Lichiardopol, N., Moree, P., Solé, P.: On Robin’s criterion for the Riemann hypothesis. *Journal de Théorie des Nombres de Bordeaux* **19**(2), 357–372 (2007). <https://doi.org/10.5802/jtnb.591>
- [3] Dusart, P.: Explicit estimates of some functions over primes. *The Ramanujan Journal* **45**, 227–251 (2018). <https://doi.org/10.1007/s11139-016-9839-4>
- [4] Hertlein, A.: Robin’s Inequality for New Families of Integers. *Integers* **18** (2018)
- [5] Mertens, F.: Ein Beitrag zur analytischen Zahlentheorie. *J. reine angew. Math.* **1874**(78), 46–62 (1874). <https://doi.org/10.1515/crll.1874.78.46>
- [6] Nazardonyavi, S., Yakubovich, S.: Superabundant numbers, their subsequences and the Riemann hypothesis. arXiv preprint arXiv:1211.2147v3 (2013), Accessed at arXiv: <https://arxiv.org/pdf/1211.2147v3.pdf>
- [7] Nicolas, J.L.: Petites valeurs de la fonction d’Euler. *Journal of Number Theory* **17**(3), 375–388 (1983). [https://doi.org/10.1016/0022-314X\(83\)90055-0](https://doi.org/10.1016/0022-314X(83)90055-0)
- [8] Nicolas, J.L., Robin, G.: Highly Composite Numbers by Srinivasa Ramanujan. *The Ramanujan Journal* **1**(2), 119–153 (1997). <https://doi.org/10.1023/A:1009764017495>
- [9] Platt, D.J., Morrill, T.: Robin’s inequality for 20-free integers. *INTEGERS: Electronic Journal of Combinatorial Number Theory* (2021)
- [10] Robin, G.: Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann. *J. Math. pures appl* **63**(2), 187–213 (1984)
- [11] Rosser, J.B., Schoenfeld, L.: Approximate Formulas for Some Functions of Prime Numbers. *Illinois Journal of Mathematics* **6**(1), 64–94 (1962). <https://doi.org/10.1215/ijm/1255631807>

Frank Vega
 NataSquad
 10 rue de la Paix
 FR 75002 Paris
 France
 e-mail: vega.frank@gmail.com