



The Riemann Hypothesis

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

May 10, 2021

The Riemann Hypothesis

Frank Vega

Abstract. Let's define $\delta(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log x - B)$, where $B \approx 0.2614972128$ is the Meissel-Mertens constant. The Robin theorem states that $\delta(x)$ changes sign infinitely often. Let's also define $S(x) = \theta(x) - x$, where $\theta(x)$ is the Chebyshev function. A theorem due to Erhard Schmidt implies that $S(x)$ changes sign infinitely often. Using the Nicolas theorem, we prove that when the inequalities $\delta(p) \leq 0$ and $S(p) \geq 0$ are satisfied for a prime $p \geq 127$, then the Riemann Hypothesis should be false. However, we could restate the Mertens second theorem as $\lim_{n \rightarrow \infty} \delta(p_n) = 0$ where p_n is the n^{th} prime number. In addition, we could modify the well-known formula $\lim_{n \rightarrow \infty} \frac{\theta(p_n)}{p_n} = 1$ as $\lim_{n \rightarrow \infty} S(p_n) = 0$. In this way, this work could mean a new step forward in the direction for finally solving the Riemann Hypothesis.

1 Introduction

In mathematics, the Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$ [1]. Let $N_n = 2 \times 3 \times 5 \times 7 \times 11 \times \dots \times p_n$ denotes a primorial number of order n such that p_n is the n^{th} prime number. Say Nicolas(p_n) holds provided

$$\prod_{q|N_n} \frac{q}{q-1} > e^\gamma \times \log \log N_n.$$

The constant $\gamma \approx 0.57721$ is the Euler-Mascheroni constant, \log is the natural logarithm, and $q | N_n$ means the prime q divides to N_n . The importance of this property is:

Theorem 1.1 [6], [7]. Nicolas(p_n) holds for all prime $p_n > 2$ if and only if the Riemann Hypothesis is true.

In mathematics, the Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

where $p \leq x$ means all the prime numbers p that are less than or equal to x . We use the following property of the Chebyshev function:

Theorem 1.2 [9]. For $x \geq 41$:

$$\theta(x) = (1 + \varepsilon(x)) \times x$$

2010 Mathematics Subject Classification: Primary 11M26; Secondary 11A41, 11A25.
Keywords: Riemann Hypothesis, Nicolas theorem, Prime numbers, Chebyshev function.

where $-\frac{1}{\log x} < \varepsilon(x) < \frac{1}{2 \times \log x}$.

Besides, in the Grönwall paper appears this:

Theorem 1.3 [3].

$$\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1.$$

Let's define $S(x) = \theta(x) - x$. Nicolas also proves that

Theorem 1.4 [7]. For $x \geq 121$:

$$\log \log \theta(x) \geq \log \log x + \frac{S(x)}{x \times \log x} - \frac{S(x)^2}{x^2 \times \log x}.$$

From the paper of Schmidt, then we can deduce that:

Theorem 1.5 [10]. $S(x)$ changes sign infinitely often.

The famous Mertens paper provides the statement:

Theorem 1.6 [5].

$$\log \left(\prod_{q \leq x} \frac{q}{q-1} \right) = \sum_{q \leq x} \frac{1}{q} + \gamma - B - \frac{1}{2} \times \sum_{q > x} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > x} \frac{1}{q^3} - \dots$$

where $B \approx 0.2614972128$ is the Meissel-Mertens constant.

Let's define:

$$\delta(x) = \left(\sum_{q \leq x} \frac{1}{q} - \log \log x - B \right),$$

Robin theorem states the following result:

Theorem 1.7 [8]. $\delta(x)$ changes sign infinitely often.

In addition, the Mertens second theorem states that:

Theorem 1.8 [5].

$$\lim_{x \rightarrow \infty} \delta(x) = 0.$$

Putting all together yields the proof that when the inequalities $\delta(p) \leq 0$ and $S(p) \geq 0$ are satisfied for a prime $p \geq 127$, then the Riemann Hypothesis should be false.

2 Central Lemma

Lemma 2.1 For a prime $p \geq 127$:

$$\frac{S(p)}{p} < 1.$$

Proof By the theorem 1.2, for all prime $p \geq 127$:

$$\begin{aligned} \frac{S(p)}{p} &= \frac{\theta(p) - p}{p} \\ &= \frac{(1 + \varepsilon(p)) \times p - p}{p} \\ &= \frac{p \times ((1 + \varepsilon(p)) - 1)}{p} \\ &= (1 + \varepsilon(p) - 1) \\ &= \varepsilon(p) \\ &< \frac{1}{2 \times \log p} \\ &< 1. \end{aligned}$$

■

3 Main Theorem

Theorem 3.1 If the inequalities $\delta(p) \leq 0$ and $S(p) \geq 0$ are satisfied for a prime $p \geq 127$, then the Riemann Hypothesis should be false.

Proof For a prime $p \geq 127$, suppose that simultaneously the inequalities Nicolas(p), $\delta(p) \leq 0$ and $S(p) \geq 0$ are satisfied. If Nicolas(p) holds, then

$$\prod_{q \leq p} \frac{q}{q-1} > e^\gamma \times \log \theta(p).$$

We apply the logarithm to the both sides of the inequality:

$$\log \left(\prod_{q \leq p} \frac{q}{q-1} \right) > \gamma + \log \log \theta(p).$$

We use that theorem 1.6:

$$\log \left(\prod_{q \leq p} \frac{q}{q-1} \right) = \sum_{q \leq p} \frac{1}{q} + \gamma - B - \frac{1}{2} \times \sum_{q > p} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > p} \frac{1}{q^3} - \dots .$$

Besides, we use that theorem 1.4:

$$\log \log \theta(p) \geq \log \log p + \frac{S(p)}{p \times \log p} - \frac{S(p)^2}{p^2 \times \log p}.$$

Putting all together yields the result:

$$\begin{aligned} & \sum_{q \leq p} \frac{1}{q} + \gamma - B - \frac{1}{2} \times \sum_{q > p} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > p} \frac{1}{q^3} - \dots \\ & > \gamma + \log \log \theta(p) \\ & \geq \gamma + \log \log p + \frac{S(p)}{p \times \log p} - \frac{S(p)^2}{p^2 \times \log p}. \end{aligned}$$

Let distribute it and remove γ from the both sides:

$$\begin{aligned} & \sum_{q \leq p} \frac{1}{q} - \log \log p - B - \frac{1}{2} \times \sum_{q > p} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > p} \frac{1}{q^3} - \dots > \\ & \frac{1}{\log p} \times \left(\frac{S(p)}{p} - \frac{S(p)^2}{p^2} \right). \end{aligned}$$

We know that $\delta(p) = \sum_{q \leq p} \frac{1}{q} - \log \log p - B$. Moreover, we know that

$$\left(\frac{S(p)}{p} - \frac{S(p)^2}{p^2} \right) \geq 0.$$

Certainly, according to the lemma 2.1, we have that $\frac{S(p)}{p} < 1$. Consequently, we obtain that $\frac{S(p)}{p} \geq \frac{S(p)^2}{p^2}$ under the assumption that $S(p) \geq 0$, since for every real number $0 \leq x < 1$, the inequality $x \geq x^2$ is always satisfied. To sum up, we would have that

$$\delta(p) - \frac{1}{2} \times \sum_{q > p} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > p} \frac{1}{q^3} - \dots > 0$$

because of

$$\frac{1}{\log p} \times \left(\frac{S(p)}{p} - \frac{S(p)^2}{p^2} \right) \geq 0.$$

However, the inequality

$$\delta(p) - \frac{1}{2} \times \sum_{q > p} \frac{1}{q^2} - \frac{1}{3} \times \sum_{q > p} \frac{1}{q^3} - \dots > 0$$

is never satisfied when $\delta(p) \leq 0$. By contraposition, $\text{Nicolas}(p)$ does not hold when $\delta(p) \leq 0$ and $S(p) \geq 0$ are satisfied for a prime $p \geq 127$. In conclusion, if $\text{Nicolas}(p)$ does not hold for a prime $p \geq 127$, then the Riemann Hypothesis should be false due to the theorem 1.1. \blacksquare

4 Discussion

The Riemann Hypothesis has been qualified as the Holy Grail of Mathematics [4]. It is one of the seven Millennium Prize Problems selected by the Clay Mathematics Institute to carry a US 1,000,000 prize for the first correct solution [2]. In the theorem 3.1, we show that if the inequalities $\delta(p) \leq 0$ and $S(p) \geq 0$ are satisfied for a prime $p \geq 127$, then the Riemann Hypothesis

should be false. Nevertheless, the well-known theorem 1.8 could be restated as

$$\lim_{n \rightarrow \infty} \delta(p_n) = 0$$

because of there are infinitely many prime numbers p_n . At the same time, we can restate the theorem 1.3 as

$$\lim_{n \rightarrow \infty} S(p_n) = 0.$$

Indeed, we think this work could help to the scientific community in the global efforts for trying to solve this outstanding and difficult problem.

References

- [1] Peter B. Borwein, Peter Borwein, Stephen Choi, Brendan Rooney, and Andrea Weirathmueller. *The Riemann Hypothesis: A Resource for the Afficionado and Virtuoso Alike*, volume 27. Springer Science & Business Media, 2008.
- [2] Keith J. Devlin. *The Millennium Problems: The Seven Greatest Unsolved Mathematical Puzzles Of Our Time*, volume 100. Basic Books New York, NY (USA), 2002.
- [3] Thomas H. Grönwall. Some asymptotic expressions in the theory of numbers. *Transactions of the American Mathematical Society*, 14(1):113–122, 1913. doi:10.2307/1988773.
- [4] Gilles Lachaud. L'hypothèse de Riemann: le Graal des mathématiciens. *La Recherche Hors-Série*, (20), 2005.
- [5] Franz Mertens. Ein Beitrag zur analytischen Zahlentheorie. *J. reine angew. Math.*, 1874(78):46–62, 1874. URL: <https://doi.org/10.1515/crll.1874.78.46>, doi:10.1515/crll.1874.78.46.
- [6] Jean-Louis Nicolas. Petites valeurs de la fonction d'Euler et hypothèse de Riemann. *Séminaire de Théorie des nombres DPP, Paris*, 82:207–218, 1981.
- [7] Jean-Louis Nicolas. Petites valeurs de la fonction d'Euler. *Journal of number theory*, 17(3):375–388, 1983. doi:10.1016/0022-314X(83)90055-0.
- [8] Guy Robin. Sur l'ordre maximum de la fonction somme des diviseurs. *Séminaire Delange-Pisot-Poitou Paris*, 82:233–242, 1981.
- [9] J. Barkley Rosser and Lowell Schoenfeld. Approximate Formulas for Some Functions of Prime Numbers. *Illinois Journal of Mathematics*, 6(1):64–94, 1962. doi:10.1215/ijm/1255631807.
- [10] Erhard Schmidt. Über die Anzahl der Primzahlen unter gegebener Grenze. *Mathematische Annalen*, 57(2):195–204, 1903.

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France
 Email: vega.frank@gmail.com